

UGT1A1

Reactivity: Human

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW: 60kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human UGT1A1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GNT1; UGT1; UDPGT; UGT1A; HUG-BR1; BILQTL1; UDPGT 1-1;

Catalog #: A6186

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 54658

Isotype: IgG

Swiss Prot: P22309

Purity: Affinity purification

For research use only.

Background:

This gene encodes a UDP-glucuronosyltransferase, an enzyme of the glucuronidation pathway that transforms small lipophilic molecules, such as steroids, bilirubin, hormones, and drugs, into water-soluble, excretable metabolites. This gene is part of a complex locus that encodes several UDP-glucuronosyltransferases. The locus includes thirteen unique alternate first exons followed by four common exons. Four of the alternate first exons are considered pseudogenes. Each of the remaining nine 5' exons may be spliced to the four common exons, resulting in nine proteins with different N-termini and identical C-termini. Each first exon encodes the substrate binding site, and is regulated by its own promoter. The preferred substrate of this enzyme is bilirubin, although it also has moderate activity with simple phenols, flavones, and C18 steroids. Mutations in this gene result in Crigler-Najjar syndromes types I and II and in Gilbert syndrome.

To place an order, please [Click HERE](#).