

CYP2C9

Reactivity: Human

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

Calculated MW: 55kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human CYP2C9

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CPC9; CYP2C; CYP2C10; CYP2C9; P450IIC9;

Catalog #: A6219

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 1559

Isotype: IgG

Swiss Prot: P11712

Purity: Affinity purification

For research use only.

Background:

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin. Studies identifying individuals who are poor metabolizers of phenytoin and tolbutamide suggest that this gene is polymorphic. The gene is located within a cluster of cytochrome P450 genes on chromosome 10q24.

To place an order, please [Click HERE](#).