

ELAC2

Reactivity:Human

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:92kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human ELAC2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ELC2; HPC2; COXPD17;

Catalog #:A7128

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:60528

Isotype:IgG

Swiss Prot:Q9BQ52

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene has a C-terminal domain with tRNA 3' trailer processing endoribonuclease activity, which catalyzes the removal of the 3' trailer from precursor tRNAs. The protein also interacts with activated Smad family member 2 (Smad2) and its nuclear partner forkhead box H1 (also known as FAST-1), and reduced expression can suppress transforming growth factor-beta induced growth arrest. Mutations in this gene result in an increased risk of prostate cancer. Multiple transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).