

IGFBP 3 Human

Description:IGFBP3 Human Recombinant produced in E.Coli is a homodimeric, non-glycosylated, polypeptide chain containing 2x264 amino acids and having a molecular mass of 28806 Dalton. IGFBP-3 is purified by proprietary chromatographic techniques.

Synonyms:Growth-hormone-dependant binding protein, IBP3, BP-53, IGFBP-3.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence:GASSGGLGPV VRCEPCDARA LAQCAPPNAV CAELVREPGC
GCCLTCLASE GQPCGIYTER CGSGLRCQPS PDEARPLQAL LDGRGLCVNA SAVSRLRAYL
LPAPPAGNA SESEEDRSAG EVESPSVSST HRVSDPKFHP LHSKIIIIKK GHAKDSQRYK
VDYESQSTDT QNFSSSESKRE TEYGPCRREM EDTLNHLKFL NVLSPRGVHI PNCDKKGFKYK
KKQCRPSKGR KR

Purity:Greater than 97.0% as determined by:(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

Formulation:

Lyophilized from a 0.2

Stability:

Lyophilized IBP3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IGF-BP 3 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Insulin-Like Growth Factor Binding Protein-3 in sterile 20mM AcOH (acetic Acid) not less than 100

Introduction:

IGFBP3 is a member of the insulin-like growth factor binding protein (IGFBP) family and encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

Biological Activity:

The ED50, calculated by its ability to inhibit IGF-II induced proliferation of MCF-7 is < 0.2

References:

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

Title: The role of insulin-like growth factor binding protein-3 in the growth inhibitory actions of androgens in LNCaP human prostate cancer cells. Publication: Article first published online: 4 OCT

2007 DOI:10.1002/ijc.23100 Copyright



Catalog #: CYPs-307

To place an order, please [Click HERE](#).

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights © 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)