

## IL 22 Mouse

**Description:** Interleukin-22 Mouse Recombinant produced in E.Coli is a single, non-glycosylated homodimeric polypeptide chain containing 147 amino acids and having a molecular mass of 16.7 kDa. The Murine IL-22 is purified by proprietary chromatographic techniques.

**Synonyms:** IL-TIF, TIFa, IL-10-related T-cell-derived-inducible factor, IL-22, ILTIF, IL-D110, zcyto18, MGC79382, MGC79384, TIFIL-23.

**Source:** Escherichia Coli.

**Physical Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Amino Acid Sequence:** The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Pro-Val-Asn.

**Purity:** Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

**Formulation:**

Lyophilized from 1X PBS.

**Stability:**

Lyophilized Interleukin-22 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL22 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

**Usage:**

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

**Solubility:**

It is recommended to reconstitute the lyophilized Interleukin -22 in sterile 18M-cm H2O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

**Introduction:**

Interleukin-22 (IL-22), also known as IL-10-related T cell-derived inducible factor (IL-TIF) was initially identified as a gene induced by IL-9 in mouse T cells and mast cells. Mouse IL-22 cDNA encodes a 179 amino acid residue protein with a putative 33 amino acids signal peptide that is cleaved to generate a 147 aa mature protein that shares approximately 79% and 22% aa sequence identity with human IL-22 and IL-10, respectively. IL-22 has been shown to activate STAT-1 and STAT-3 in several hepatoma cell lines and upregulate the production of acute phase proteins. IL-22 is produced by normal mouse T cells upon Con A activation. Mouse IL-22 expression is also induced in various organs upon lipopolysaccharide injection, suggesting that IL-22 may be involved in inflammatory responses. The functional IL-22 receptor complex consists of two receptor subunits, IL-22R (previously an orphan receptor named CRF2-9) and IL-10R (previously known as CRF2-4), belonging to the class II cytokine receptor family.

**Biological Activity:**

The ED50 as determined by its ability to induce IL-10 secretion in Colo205 cells is less than

www.neobiolab.com  
info@neobiolab.com  
888.754.5670, +1 617.500.7103 United States  
0800.088.5164, +44 020.8123.1558 United Kingdom  
0.5ng/ml, corresponding to a Specific Activity of 2,000,000IU/mg.



---

***To place an order, please [Click HERE](#).***

Catalog #:CYP5-546

For research use only.

© 2015 NeoScientific

