www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom

CD244 Human

Catalog #:PRPS-1180

For research use only.

Description:CD244 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 230 amino acids (19-224 a.a) and having a molecular mass of 25.5kDa.CD244 is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Synonyms: Natural killer cell receptor 2B4, NK cell activation-inducing ligand, NAIL, NK cell type I receptor protein 2B4, NKR2B4, h2B4, CD244, 2B4, Nmrk, SLAMF4.

Source:E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MGSHGKGCQG SADHVVSISG VPLQLQPNSI QTKVDSIAWK KLLPSQNGFH HILKWENGSL PSNTSNDRFS FIVKNLSLLI KAAQQQDSGL YCLEVTSISG KVQTATFQVF VFDKVEKPRL QGQGKILDRG RCQVALSCLV SRDGNVSYAW YRGSKLIQTA GNLTYLDEEV DINGTHTYTC NVSNPVSWES HTLNLTQDCQ NAHQEFRFWP.

Purity: Greater than 85% as determined by SDS-PAGE.

Formulation:
CD244 protein solution (0.5mg/ml) containing 20mM Tris-HCl buffer, pH8.0 and 0.4M Urea.

Stability:
Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:
NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drµgs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:
Cluster of Differentiation 244 (CD244/Natural Killer Cell Receptor 2B4) contains 2 Ig-like (immunoglobulin-like) domains. CD244 is a cell surface receptor expressed on natural killer (NK) cells (and some T cells) which mediate non-major histocompatibility complex (MHC) restricted killing. NK-cell and target cells interaction via the CD244 receptor is believed to modulate NK-cell cytolytic activity. CD244 molecules function in eosinophil adhesion and chemotaxis, and have a role in the pathophysiology of allergic rhinitis (AR).

To place an order, please Click HERE.