

CD3G Human

Description: CD3G Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 117 amino acids (23-116a.a.) and having a molecular mass of 13.1kDa. CD3G is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PRPS-1399

For research use only.

Synonyms: T-cell surface glycoprotein CD3 gamma chain, CD3-GAMMA, T3G, CD3G.

Source: E.coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHH SSGLVPRGSH MGSQSIKGNH LVKVYDYQED
GSVLLTCDAE AKNITWFKDG KMIGFLTEDK KKWNLGSNAK DPRGMYQCKG SQNKSKPLQV
YYRMCQCIE LNAATIS.

Purity: Greater than 90% as determined by SDS-PAGE.

Formulation:

CD3G protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.4M urea and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Short chain dehydrogenase/reductase family 16C, member5 (SDR16C5) is active in the oxidative direction as well as in the reductive one. SDR16C5 oxidizes all-trans-retinol in all-trans-retinaldehyde. No activity was detected with 11-cis-retinol or 11-cis-retinaldehyde as substrates with either NAD⁺/NADH or NADP⁺/NADPH.

To place an order, please [Click HERE](#).